


Example: Computer chips have a 25% chance of being defective. Create the probability distribution for the # of defective chips in a sample of 3.



x	0	1	2	3
P(x)	0.422	0.422	0.141	0.016

CH. 17 PROBABILITY MODELS

Binomial models:

* Interested in the number of successes in a set # of trials $n=10$

* 4 conditions that must apply:

- Only 2 possible outcomes (success/failure) *right* *wrong*

- Probability of success remains constant (called p) $p=0.25$

- Number of trials is set/known (called n) $n=10$

- Independent trials

$n=3$
 $pop \geq 30$ * **10% condition:** If we cannot assume independence we can proceed as long as the sample is smaller than 10% of the population $(pop \geq 10n)$

* If these 4 conditions apply, we have a **Bernoulli trial**

Notation:

$B(n, p)$

$P(X=k)$

prob. of success

$$\mu_X = n \cdot p$$

$$\text{Ex: } 10 \times 0.25 = 2.5$$

$$\sigma_X = \sqrt{n \cdot p \cdot (1-p)}$$

$$\text{Ex: } \sqrt{10 \cdot 0.25 \cdot 0.75} =$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A|B)$$

Binomial probabilities:

Example (from earlier):

$$p=0.25 \quad n=3$$

Computer chips have a 25% chance of being defective. Create the probability distribution for the # of defective chips in a sample of 3.

(a) What is the probability of having 2 or more defective chips?

(b) What is the probability of having 1 or less defective chips?

(c) What is the probability of having exactly 2 defective chips?

STEP 1: Bernoulli? check to see....

① success = defective
failure = non-defective

② $p=0.25$ + constant

③ $n=3$

④ assume indep.

all comp chips ≥ 30 ? yes.

$$B(3, 0.25)$$

STEP 2: Create probability distribution (we did this before)

X	0	1	2	3
P(X)	0.42188	0.42188	0.14063	0.01563

STEP 3: ANSWER THE QUESTIONS (notation!!)

(a) What is the probability of having 2 or more defective chips?

$$P(X \geq 2) = 0.15626$$

(b) What is the probability of having 1 or less defective chips?

$$P(X \leq 1) = 0.84376$$

(c) What is the probability of having exactly 2 defective chips?

$$P(X = 2) = 0.14063$$

$$\mu_x =$$
$$\sigma_x =$$

QUICKER WAY TO GET PROBABILITIES:

Formula:

$$P(X = k) = ({}_nC_k)(p^k)(q^{n-k})$$

combination: ${}_nC_r$

$1 - p = \text{prob. failure}$

Same example: $B(3, 0.25)$

X	0	1	2	3
P(X)	0.421875	0.421875	0.140625	0.015625

$$P(X=0) = ({}_3C_0)(0.25^0)(0.75^3)$$
$$({}_3nCr_0)(0.25^1)(0.75^2)$$

Example: I am playing a game in which I have only a 39% chance of winning. I am playing 4 times. Create the probability distribution below: $P(X=k) = {}_n C_k (p^k)(q^{n-k})$

$X(k)$	$P(X) \quad P(X=k)$	$B(4, 0.39)$
0	$({}_4 C_0)(0.39^0)(0.61^4) = 0.13845841$	
1	$({}_4 C_1)(0.39^1)(0.61^3) = 0.35409036$	
2	$({}_4 C_2)(0.39^2)(0.61^2) = 0.33957816$	
3	$({}_4 C_3)(0.39^3)(0.61^1) = 0.14473836$	
4	$({}_4 C_4)(0.39^4)(0.61^0) = 0.02313441$	

X	P(X)
0	0.13846
1	0.35409
2	0.33958
3	0.14474
4	0.02313

B(4, 0.39)

So let's answer some easy questions:

$$P(X=2) =$$

$$P(X < 2) =$$

$$P(X \geq 3) =$$

$$P(2 \leq X \leq 4) =$$

$$\mu_x = n \cdot p$$

$$\sigma_x = \sqrt{n \cdot p \cdot q}$$

Now let's change the sample size to 10. $B(10, 0.39)$

x	$P(x)$
0	
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	

$$P(X=9) =$$

$$P(X < 4) =$$

$$P(X \geq 6) =$$

$$P(5 \leq X \leq 7) =$$

$$n = 100$$

$$P(30 \leq X \leq 50)$$

Would you want to answer these questions for a sample size of 50? Of 100? NO! **So we can use the calculator!**

For **$P(X=k)$**

- Use ... `binompdf(n, p, k)`

2nd Distrib → O/A

X				
P(X)				

- k = the number you are looking for...

Example: $P(X = 5)$ $k = 5$

- pdf = probability distribution function
(gives each individual outcome's probability)

For $P(X \leq k)$

- Use ... **binomcdf**(n, p, k)

$$P(X < 10)$$

- k = the number you are looking for... $P(X \leq 9)$

Example: $P(X \leq 6)$ k = 6

$$P(X > 7)$$

$$1 - P(X \leq 7)$$

- Notice that it ONLY GIVES YOU: LESS THAN OR
EQUAL TO

- cdf = cumulative distribution function... adds up all the probabilities below

$$\text{Ex: } P(X \leq 3) = P(X=0) + P(X=1) + P(X=2) + P(X=3)$$

HW answers:

- 1)
 - (a) Yes
 - (b) Yes
 - (c) No- no "success"
 - (d) No- more than 2 outcomes
 - (e) No- more than 2 outcomes
 - (f) No- probability of success isn't constant
 - (g) Yes
 - (h) Yes
 - (i) No- more than 2 outcomes
 - (j) Yes

2) Binomial?

- succes = alarm failing failure = alarm working
- $p = 0.05$ and is constant
- Independent trials stated
- $n = 6$

$B(6, 0.05)$

(a) $P(X = 3) = \text{binompdf}(6, 0.05, 3) = 0.002$

(b) $P(X < 2) = P(X \leq 1) = \text{binomcdf}(6, 0.05, 1) = 0.9672$

(c) $P(X = 0) = \text{binompdf}(6, 0.05, 0) = 0.7351$

Example: John is taking archery. He has a 30% chance of hitting the target each time he shoots. He shoots 8 times. (Lets do this example together)

1) What is the probability that he hits the target 4 times? $B(8, 0.30)$
 $P(X=4) = \text{binompdf}(8, 0.3, 4) = 0.1361$

2) What is the probability that he hits the target 2 times or less?
 $P(X \leq 2) = \text{binomcdf}(8, 0.3, 2) = 0.5518$

3) What is the probability that he hits the target at least 3 times?
 $P(X \geq 3) = 1 - P(X \leq 2) = 1 - \text{binomcdf}(8, 0.3, 2) = 0.4482$

4) What is the probability that he hits the target less than 5 times?
 $P(X < 5) = P(X \leq 4) = \text{binomcdf}(8, 0.3, 4) = 0.942$

5) What is the probability that he hits the target more than 6 times?
 $P(X > 6) = 1 - P(X \leq 6) = 1 - \text{binomcdf}(8, 0.3, 6) = 0.0013$

6) How many times do we expect him to hit the target? (average!)
 $\mu_x = n \cdot p = 8 \cdot 0.3 = 2.4 \text{ hits}$

7) What is the standard deviation of the number of times he hits the target?
 $\sigma_x = \sqrt{n \cdot p \cdot q} = \sqrt{(8)(0.3)(0.7)} = 1.296 \text{ hits}$

Example: John is taking a target class and has a 30% chance of hitting the target each time he shoots. ***ANSWERS*** $B(8, 0.30)$

1) What is the probability that he hits the target 4 times?

$$P(X = 4) = \text{binompdf}(8, 0.30, 4) =$$

2) What is the probability that he hits the target 2 times or less?

$$P(X \leq 2) = \text{binomcdf}(8, 0.30, 2) =$$

3) What is the probability that he hits the target at least 3 times?

$$P(X \geq 3) = 1 - P(X \leq 2) = 1 - \text{binomcdf}(8, 0.30, 2) =$$

4) What is the probability that he hits the target less than 5 times?

$$P(X < 5) = P(X \leq 4) = \text{binomcdf}(8, 0.30, 4) =$$

5) What is the probability that he hits the target more than 6 times?

$$P(X > 6) = P(X \geq 7) = 1 - P(X \leq 6) = 1 - \text{binomcdf}(8, 0.30, 6) =$$

6) How many times do we expect him to hit the target? (average!)

$$E(X) = \mu_X = 8 * 0.30 = 2.4 \text{ times}$$


7) What is the standard deviation of the number of times he hits the target?

$$\sigma_X = \text{sqrt}(8 * 0.30 * 0.70) = 1.296 \text{ times}$$

Try this example on your own: 150 businesses are sent mailings asking them to answer a survey question and send the mailing back. The probability of nonresponse is 55%.

$$B(150, 0.45)$$

- 1) What is the average number of businesses that WILL respond?
- 2) What is the std. deviation of the number that will respond?
- 3) What is the probability that 75 businesses will respond?
- 4) What is the probability that 60 businesses or less will respond?
- 5) What is the probability that 60 businesses or more will respond?
- 6) What is the probability that less than 60 businesses will respond?
- 7) What is the probability that greater than 60 businesses will respond?
- 8) What number of surveys would you have to send out if you wanted to be able to expect to get 90 back?
- 9) What is the probability that between 50 and 70 businesses will respond?



$$P(51 \leq X \leq 69)$$

$$P(50 < X < 70)$$

1) $\mu_X = 67.5$ } businesses

2) $\sigma_X = 6.093$

3) $P(X = 75) = 0.0306$ PDF

4) $P(X \leq 60) = 0.1251$ CDF

5) $P(X \geq 60) = 1 - P(X \leq 59) = 0.9058$

6) $P(X < 60) = P(X \leq 59) = 0.0942$

7) $P(X > 60) = 1 - P(X \leq 60) = 0.8749$

8) $n = 200$

9) $P(50 < X < 70) = P(51 \leq X \leq 69)$

$$= P(X \leq 69) - P(X \leq 50)$$

$$= 0.6271$$

$$B(150, 0.45)$$

$$\mu_X = n \cdot p$$
$$90 = n \cdot 0.45$$

Complete worksheet from yesterday

#3 -- 8

$$\textcircled{3} B(10, 0.25) \quad P(X \geq 6) = 1 - P(X \leq 5) = 0.0197$$

$$\textcircled{4} B(15, 0.90) \quad P(X \geq 9) = 1 - P(X \leq 8) = 0.9997$$

$$\textcircled{5} B(10, 0.60) \quad P(X = 3) = 0.0425$$

$$\textcircled{6} B(9, 0.60) \quad \textcircled{a} P(X \geq 5) = 1 - P(X \leq 4) = 0.7334$$

$$\textcircled{b} P(X = 7) = 0.1612$$

$$\textcircled{c} P(X > 3) = 1 - P(X \leq 3) = 0.9006$$

⑧ $\mu \pm \sigma$

(160.6878, 171.3122)



$$P(161 \leq X \leq 171)$$


$$P(X \leq 171) - P(X \leq 160) = 0.6998$$

$$\text{binomcdf}(200, 0.83, 171) - \text{binomcdf}(200, 0.83, 160)$$

Complete the experiment in the notes


(If you dont have the notes, let me know- I have extra copies of the expt)

x	$P(x)$
0	0.17851
1	0.38448
2	0.31054
3	0.11148
4	0.01501



Complete the experiment

$n = 4$




$p = 0.35$


$p = 0.65$




$n = 10$



$n = 20$



$n = 30$



N

Ch. 17: Probability Models:

Binomial Random Variables: LARGE SAMPLE SIZE

What happens to the shape of the Binomial Random Variable when n is large?

Skew \longrightarrow normal
(small n) (large n)

What is considered a "large enough" n (for the shape to look normal)??

check:

$$\begin{matrix} n \cdot p \\ n \cdot q \end{matrix} \geq 10$$

then n is
large enough
to use
 $N(,)$

So if the check passes...

- We can say that the distribution is approx. normal,

and can use normalcdf!

$$B(n, p)$$

$$n \cdot p \geq 10$$

- Calculator:

normalcdf(lower bound, upper bound, mean, std. dev) $N()$

$$P(X=30) = \text{PDF}$$

- Same mean and std dev. that we learned before:

$$\mu_X = n \cdot p$$

$$\sigma_X = \sqrt{n \cdot p \cdot (1-p)}$$

Example 1:

It is said that 75% of people pay their credit card bill on time. If we take a sample of 125 adults, what is the chance that over 80 of them paid their bill on time this past month?

Bernoulli?

- succ/fail
- p
- n
- indep.

$$B(125, 0.75)$$

Check:

$$125 \cdot 0.75 \neq 10$$
$$125 \cdot 0.25 \neq 10$$
$$N(93.75, 4.8412)$$

Work:

$$P(X \geq 80) = 0.9977$$

$$\text{normcdf}(80, \infty, 93.75, 4.8412)$$

Example 2:

Assume 15% of adults jog regularly. We survey 1,000 adults what is the probability that between 120 and 160 people in our sample jogs?

$$B(1000, 0.15)$$

Check:

$$\begin{array}{l} 1000 \cdot 0.15 \\ 1000 \cdot 0.85 \end{array} \approx 10$$

$$N(150, 11.2916)$$

$$P(120 < X < 160) =$$

$$0.8081$$

Complete worksheet #1 -- 4, 10

Answers:

1) (a) yes (b) no (c) no

2) (a) $n = 100$ (b) $n = 34$ (c) $n = 50$

3) $B(700, 0.05)$ Check--> passes $(700)(0.05) \geq 10$
 $(700)(0.95)$

$$P(X > 50) = 0.0046$$

$N(35, 5.7663)$

4) $B(400, 0.48)$ Check --> passes $(400)(0.48) \geq 10$
 $(400)(0.52)$

$\nwarrow 0.55 \times 400$

$$P(180 < X < 220) = 0.8826$$

$\nearrow 0.45 \times 400$

$N(192, 9.992)$

10) $B(2000, 0.5)$ check --> passes $(2000)(0.5) \geq 10$
 $(2000)(0.5)$ ✓

$$P(975 < X < 1050) = 0.8556$$

$$N(1000, 22.361)$$

$$\sqrt{(2000)(0.5)(0.5)}$$

p. 402

#18) B(6, 0.80)

$$(a) P(B^c \cap B^c \cap B) = (0.20)(0.20)(0.80) = \mathbf{0.032}$$

(b) $B(6, 0.20)$ $p = 0.20$ because we are concerned with MISSING

$$P(X \geq 1) = 1 - P(X \leq 0) = \text{binomcdf}(6, 0.20, 0) = \mathbf{0.7379}$$

$$(c) P(B^c \cap B^c \cap B^c \cap B) = 0.0064$$

$$P(B^c \cap B^c \cap B^c \cap B^c \cap B) = 0.00128 \quad \text{red } > \quad = \mathbf{0.00768}$$

$$(d) P(X = 4) = \text{binompdf}(6, 0.80, 4) = \mathbf{0.24576}$$

$$(e) P(X \geq 4) = 1 - P(X \leq 3) = 1 - \text{binomcdf}(6, 0.80, 3) = \mathbf{0.90112}$$

$$(f) P(X \leq 4) = \text{binomcdf}(6, 0.80, 4) = \mathbf{0.34464}$$

29) B(300, 0.06)

(a) check: $(300)(0.06)$
 $(300)(0.94) \geq 10 \quad \rightarrow N(18, 4.113)$

(b) $P(X < 12) = \text{normalcdf}(-E99, 12, 18, 4.113) = \mathbf{0.0723}$

(c) $P(X > 50) = \text{normalcdf}(50, E99, 18, 4.113) = \mathbf{3.654 \times 10^{-15}}$
No, it is not likely.

$$\sqrt{(300)(0.06)(0.94)}$$

$$\mu_x = n \cdot p$$

$$\sigma_x = \sqrt{n \cdot p \cdot (1-p)}$$

On the worksheet:
#5, 6, 9

5) $B(100, 0.70)$

Check:

$$100 \cdot 0.70 \geq 10$$

$$100 \cdot 0.30$$

$N(70, 4.5826)$

$$P(X > 80) = 0.0145$$

6) $B(1500, 0.56)$

Check:

$$1500 \cdot 0.56 \geq 10$$

$$1500 \cdot 0.44$$

$N(840, 19.225)$

$$P(X > 750) = 0.9999$$

9) $B(1000, 0.10)$

not

0.9 = does


Check:

$$1000 \cdot 0.90 \geq 10$$

$$1000 \cdot 0.10$$

$N(100, 9.4868)$

$$P(X < 100) = 50\%$$



Book problems:


p. 402 #17 & 21, ~~25 & 30~~ *26 & 30*

17) B(5, 0.13)

(a) $P(L^c \cap L^c \cap L^c \cap L^c \cap L) = (0.87) (0.87) (0.87) (0.87)(0.13) = \mathbf{0.07448}$

(b) $P(X \geq 1) = 1 - P(X \leq 0) = 1 - \text{binomcdf}(5, 0.13, 0) = \mathbf{0.5016}$

(c) second: $P(L^c \cap L) = 0.1131$

third: $P(L^c \cap L^c \cap L) = 0.0984$  $= \mathbf{0.2115}$

(d) $P(X = 3) = \text{binompdf}(5, 0.13, 3) = \mathbf{0.0166}$

(e) $P(X \geq 3) = 1 - P(X \leq 2) = 1 - \text{binomcdf}(5, 0.13, 2) = \mathbf{0.0179}$

(f) $P(X \leq 3) = \text{binomcdf}(5, 0.13, 3) = \mathbf{0.9987}$

21) B(12, 0.87)

(a) $\mu_X = 12 \cdot 0.87 = \mathbf{10.44 \text{ people}}$

$$\sigma_X = \sqrt{12 \cdot 0.87 \cdot 0.13} = \mathbf{1.165 \text{ people}}$$

(b) (i) $1 - P(X = 12) = 1 - \text{binompdf}(12, 0.87, 12) = \mathbf{0.8196}$

(ii) $P(X \leq 10) = \text{binomcdf}(12, 0.87, 10) = \mathbf{0.4748}$

(iii) $P(X = 6) = \text{binompdf}(12, 0.87, 6) = \mathbf{0.0019}$

(iv) $P(X > 6) = 1 - P(X \leq 6) = 1 - \text{binomcdf}(12, 0.87, 6) = \mathbf{0.9978}$

26) B(12, 0.125)

(a) $P(X = 0) = \text{binompdf}(12, 0.125, 0) = \mathbf{0.2014}$

(b) $P(X \geq 2) = 1 - P(X \leq 1) = 1 - \text{binomcdf}(12, 0.125, 1) = \mathbf{0.4533}$

(c) $P(X = 3) + P(X = 4) =$
 $= \text{binompdf}(12, 0.125, 3) + \text{binompdf}(12, 0.125, 4) = \mathbf{0.1707}$

(d) $P(X \leq 4) = \text{binomcdf}(12, 0.125, 4) = \mathbf{0.9887}$

30) B(150, 0.125)

(a) $\mu_X = 18.75$ frogs

$\sigma_X = 4.05$ frogs

(b) check: $\begin{matrix} 150 \cdot 0.125 \\ 150 \cdot 0.875 \end{matrix} \geq 10 \quad N(18.75, 4.05)$

(c) $P(X > 22) = 0.211$

No

Ch. 17 Classwork